

LETTERS TO THE EDITOR

Symbiotic Circulation

There have been excellent reviews of the pathways, distribution and functional significance of the coronary collateral circulation (1-4). However, none of the reviews has described a unique pattern of *symbiotic circulation* which occurs when two diseased vessels each supply collateral circulation to the other. This collateral *symbiotic circulation* pattern is present in an estimated 1% of the 950 coronary arteriograms performed annually in our laboratory.

An example of this phenomenon is shown in Figure 1. In panel A there is injection of contrast material into the right coronary artery in the left anterior oblique projection. There is complete occlusion of the distal right coronary artery and no filling of the posterior descending branch. This diseased right coronary artery provides collateral channels to the left anterior descending artery (arrows) via the conus branch of the right coronary artery. Panel B is a right anterior oblique projection of the left coronary injection. The left anterior descending artery is occluded proximally. However, this left system provides collateral channels to the posterior descending branch (arrows) of the right coronary artery by way of collateral vessels from the obtuse marginal branches of the left circumflex artery. There are also collateral vessels from the diagonal branches to the distal left anterior descending artery.

Other patterns of double coronary collateral circulation are possible, but when two diseased arteries are in mutual dependence, it may be called a *symbiotic circulation*.

Arthur Dodek, MD, FACC
Director, Cardiac Catheterization Laboratory
St. Paul's Hospital
University of British Columbia
Vancouver, British Columbia, Canada

Address for correspondence:
#503 - 1160 Burrard St.
Vancouver, British Columbia, Canada V6Z2E8

References

1. Gensini GG, Da Costa BCB. The coronary collateral circulation in living man. *Am J Cardiol* 1969;24:393-400.
2. James TN. The delivery and distribution of coronary collateral circulation. *Chest* 1970;58:183-203.
3. Helfant RH, Vokonas PS, Gorlin R. Functional importance of the human coronary collateral circulation. *N Engl J Med* 1971;284:1277-81.
4. Levin DC. Pathways and functional significance of the coronary collateral circulation. *Circulation* 1974;50:831-9.

Comment

Dr. Dodek's "symbiotic circulation" is an elegant term for a phenomenon commonly observed in the coronary arteriograms of





Figure 1. Coronary arteriograms demonstrating pattern of symbiotic circulation. See text.