

**\*Daniel C. Link, MD**  
**Richard G. Bach, MD**

\*Washington University  
Division of Oncology  
660 South Euclid  
Campus Box 8007  
Saint Louis, Missouri 63110  
E-mail: dlink@im.wustl.edu

doi:10.1016/j.jacc.2007.01.055

## REFERENCES

- Güven H, Shepherd RM, Bach RG, Capoccia BJ, Link DC. The number of endothelial progenitor cell colonies in the blood is increased in patients with angiographically significant coronary artery disease. *J Am Coll Cardiol* 2006;48:1579-87.
- Leor J, Marber M. Endothelial progenitors: a new Tower of Babel? *J Am Coll Cardiol* 2006;48:1588-90.
- Yoder MC, Mead LE, Prater D, et al. Re-defining endothelial progenitor cells via clonal analysis and hematopoietic stem/progenitor cell principals. *Blood* 2006 Oct 19; [E-pub ahead of print].
- George J, Goldstein E, Abashidze S, et al. Circulating endothelial progenitor cells in patients with unstable angina: association with systemic inflammation. *Eur Heart J* 2004;25:1003-8.
- Sandri M, Adams V, Gielen S, et al. Effects of exercise and ischemia on mobilization and functional activation of blood-derived progenitor cells in patients with ischemic syndromes: results of 3 randomized studies. *Circulation* 2005;111:3391-9.

## Adjunctive Thrombectomy in Acute Myocardial Infarction: For Some but Not for All

We read with interest the report by Silva-Orrego et al. (1) regarding the effects of thrombus aspiration on myocardial reperfusion in the setting of primary angioplasty. The study reported a significant enhancement in myocardial reperfusion by a strategy of manual thrombus aspiration before stenting on top of abciximab administration. However, in our opinion some concerns exist about the results of this study. First, the investigators randomized patients with acute myocardial infarction (MI) to a conventional stent-assisted primary angioplasty versus a strategy applying manual aspiration device before stent, regardless of angiographic appearance of thrombus. Early studies evaluating the feasibility of thrombectomy in the setting of thrombus-containing lesion and acute MI clearly reported on the efficacy of thrombectomy devices in reducing thrombus burden at target lesion, consequently limiting distal embolization and no-reflow phenomenon (2,3).

Unfortunately, Silva-Orrego et al. (1) did not provide data concerning thrombus burden at baseline angiography, nor on thrombus removal after device use. Thus, the reported reduction in distal embolization rate may not be univocally ascribed to the device effect, as other factors may affect this result, such as differences in thrombus burden of culprit lesions.

Second, the researchers found a rate of postprocedural myocardial blush grade (MBG) 2/3 in both arms, which is surprisingly higher than expected, based on Thrombolysis In Myocardial Infarction (TIMI) flow achieved after the intervention, and on

time-to-treatment reported in this cohort. In the largest study validating the angiographic reperfusion, MBG was strongly affected by epicardial flow, being that patients showing TIMI flow grade <3 very rarely reached an adequate blush (4). Indeed, it has been shown that the effectiveness of myocardial reperfusion after primary angioplasty, as assessed by ST-segment resolution and MBG is clearly time-dependent (5).

In conclusion, to avoid conflicting results, we believe that future studies evaluating thrombectomy as adjunctive strategy in the setting of primary angioplasty should enroll only patients with angiographically documented thrombus at the lesion site, and that reperfusion findings should be carefully evaluated according to time-to-treatment.

**\*Massimo Napodano, MD**  
**Angelo Ramondo, MD**  
**Sabino Iliceto, MD, FACC**

\*Department of Cardiac, Thoracic, and Vascular Sciences  
University of Padova  
2 via Giustiniani  
35128 Padova  
Italy  
E-mail: massimo.napodano@sanita.padova.it

doi:10.1016/j.jacc.2007.01.056

## REFERENCES

- Silva-Orrego P, Colombo P, Bigi R, et al. Thrombus aspiration before primary angioplasty improves myocardial reperfusion in acute myocardial infarction: the DEAR-MI study. *J Am Coll Cardiol* 2006;48:1552-9.
- Silva JA, Ramee SR, Choen DJ, et al. Rheolytic thrombectomy during percutaneous revascularization for acute myocardial infarction: experience with the Angiojet catheter. *Am Heart J* 2001;141:353-9.
- Napodano M, Pasquetto G, Saccà S, et al. Intracoronary thrombectomy improves myocardial reperfusion during direct angioplasty for acute myocardial infarction. *J Am Coll Cardiol* 2003;42:1395-402.
- van't Hof AWJ, Liem A, Suryapranata H, Horntje JCA, de Boer MJ, Zijlstra F on behalf of the Zwolle Myocardial Infarction Study Group. Angiographic assessment of myocardial reperfusion in patients treated with primary angioplasty for acute myocardial infarction: myocardial blush grade. *Circulation* 1998;97:2302-6.
- De Luca G, van't Hof AW, de Boer MJ, et al. Time-to-treatment significantly affects the extent of ST-segment resolution and myocardial blush in patients with acute myocardial infarction treated by primary angioplasty. *Eur Heart J* 2004;25:1009-13.

## Reply

We appreciated the comments by Dr. Napodano and colleagues and would like to address their concern. The DEAR-MI (De-thrombosis to Enhance Acute Reperfusion in Myocardial Infarction) study (1) included consecutive randomization of acute myocardial infarction (AMI) patients fulfilling the inclusion criteria regardless of angiographic evidence of thrombus based on the assumption that thrombus is always present in AMI. Compared with angiography, the presence of intracoronary thrombus is significantly underestimated by angiography (2). In our randomized population, 73% of controls and 81% of patients undergoing aspiration showed an occluded vessel with Thrombolysis In Myocardial Infarction (TIMI) flow grade 0/1 at the initial angiogram,