

QUARTERLY FOCUS ISSUE: PREVENTION/OUTCOMES

Alcohol Consumption and Cardiovascular Mortality Among U.S. Adults, 1987 to 2002

Kenneth J. Mukamal, MD, MPH, MA,* Chung M. Chen, MA,† Sowmya R. Rao, PhD,‡
Rosalind A. Breslow, PhD¶

Boston, Massachusetts; and Rockville, Maryland

- Objectives** The aim of this study was to determine the association of alcohol consumption and cardiovascular mortality in the U.S. population.
- Background** Alcohol consumption has been associated with a lower risk of cardiovascular disease in cohort studies, but this association has not been prospectively examined in large, detailed, representative samples of the U.S. population.
- Methods** We analyzed 9 iterations of the National Health Interview Survey, an annual survey of a nationally representative sample of U.S. adults between 1987 and 2000. Exposures of interest included usual volume, frequency, and quantity of alcohol consumption and binge drinking. Mortality was ascertained through linkage to the National Death Index through 2002. Relative risks were derived from random-effects meta-analyses of weighted, multivariable-adjusted hazard ratios for cardiovascular mortality from individual survey administrations.
- Results** Light and moderate volumes of alcohol consumption were inversely associated with cardiovascular mortality. Compared with lifetime abstainers, summary relative risks were 0.95 (95% confidence interval [CI]: 0.88 to 1.02) among lifetime infrequent drinkers, 1.02 (95% CI: 0.94 to 1.11) among former drinkers, 0.69 (95% CI: 0.59 to 0.82) among light drinkers, 0.62 (95% CI: 0.50 to 0.77) among moderate drinkers, and 0.95 (95% CI: 0.82 to 1.10) among heavy drinkers. The magnitude of lower risk was similar in subgroups of sex, age, or baseline health status. There was no simple relation of drinking pattern with risk, but risk was consistently higher among those who consumed ≥ 3 compared with 2 drinks/drinking day.
- Conclusions** In 9 nationally representative samples of U.S. adults, light and moderate alcohol consumption were inversely associated with CVD mortality, even when compared with lifetime abstainers, but consumption above recommended limits was not. (J Am Coll Cardiol 2010;55:1328–35) © 2010 by the American College of Cardiology Foundation

Alcohol consumption has been consistently associated with a lower risk of cardiovascular disease (CVD) in epidemiological studies (1,2), an association attributed in great part to the increase in high-density lipoprotein cholesterol (HDL-C) caused by alcohol consumption (3).

However, a number of uncertainties about the association of alcohol consumption and CVD remain, punctuated by

the absence of a long-term randomized controlled trial on CVD events with which to confirm the results of observational studies. These uncertainties include potentially diverse effects on coronary heart disease (CHD) and stroke (4), inclusion of former or occasional drinkers with long-term abstainers as a referent category (5), generalizability to the adult U.S. population (6), and the importance of

See page 1336

From the *Division of General Medicine and Primary Care, Beth Israel Deaconess Medical Center, Boston, Massachusetts; †CSR Incorporated, Arlington, Virginia; ‡Biostatistics Center and Institute for Health Policy, Massachusetts General Hospital, Boston, Massachusetts; and the ¶Division of Epidemiology and Prevention Research, National Institute on Alcohol Abuse and Alcoholism, Rockville, Maryland. Computer programming and statistical support were provided through the Alcohol Epidemiologic Data System funded by contracts N0AA32007 and HHSN267200800023C from the National Institute on Alcohol Abuse and Alcoholism (NIAAA). The NIAAA reviewed and approved this report before submission. The findings and conclusions in this report are those of the authors and not necessarily those of the agency. Dr. Rao has received funding from GE Corporate Healthcare.

Manuscript received August 4, 2009; revised manuscript received September 16, 2009, accepted October 14, 2009.

drinking patterns in modifying the association (7). Measures of overall volume of alcohol consumption obscure the relative contributions of drinking frequency (how often alcohol is consumed), drinking quantity (how much alcohol is typically consumed on those days), and binge drinking (episodes of 5 or more drinks in a day); and their individual contributions to CVD risk have not been thoroughly investigated.

To evaluate the associations of alcohol consumption and drinking patterns with CVD, cerebrovascular, and

CHD mortality in a nationally representative sample of U.S. adults, we aggregated data from 9 administrations of the National Health Interview Survey (NHIS). The NHIS is an annual in-person survey conducted by the National Center for Health Statistics (NCHS) of the Centers for Disease Control and Prevention and has included detailed questions on alcohol consumption on selected surveys since 1987.

Methods

Study sample. Since 1957, the NCHS has conducted annual cross-sectional household interview surveys of the noninstitutionalized civilian U.S. population. The NHIS data are collected during a personal household interview conducted by interviewers employed and trained by the U.S. Bureau of the Census. The sampling plan follows a multi-stage area probability design, with oversampling of selected minority populations, including both black and Hispanic persons. Survey participation is voluntary, and the confidentiality of responses is assured by the Public Health Service Act.

The content of the survey is updated approximately every 10 to 15 years and has remained relatively unchanged since 1997. The NHIS questionnaire from 1982 to 1996 included a core set of basic health and demographic items and 1 or more added sets of questions on diverse health topics but did not consistently include questions about health-related behaviors, such as alcohol consumption. Since 1997, the NHIS questionnaire has similar core questions that include demographic information, health status and limitations, health insurance, income and assets, and diverse health behaviors.

Our study used data from NHIS years that included alcohol consumption data (1987 to 1988, 1990 to 1992, and 1997 to 2000). Response rates for alcohol-related modules for adults included in these NHIS years range from 70% in 1999 to 87% in 1988. All NHIS data are publicly available at the NCHS website.

Alcohol consumption. As noted, alcohol consumption was ascertained variably in the NHIS. Table 1 summarizes the methods used. In 1987 and 1992, a validated food

frequency questionnaire (8) was administered that included separate quantity-frequency items for beer, wine, and liquor. In 1988 and 1997 to 2000, alcohol-related items included: 1) intake of 12 or more drinks in one's lifetime, in any previous year, and in the past year; and 2) drinking frequency, drinking quantity, and binge drinking in the past year. In 1990, similar questions were asked but with a time frame of the last 2 weeks for current drinking; only individuals who reported that those 2 weeks were typical for consumption in the past year were included in these analyses. Quantity-frequency questions were also administered in 1991 as part of a Drug and Alcohol Use module to adults age 18 to 44 years, although intake in any previous year was not queried. Binge drinking was ascertained in 1988, 1991, and 1997 to 2000.

From these questions, we derived 6 alcohol consumption categories. Current abstainers were categorized as never drinkers (<12 drinks in one's lifetime), lifetime infrequent drinkers (>12 drinks in one's lifetime, <12 drinks in any previous year), and former drinkers (>12 drinks in one's lifetime, >12 drinks in a previous year). Current drinkers were categorized as light (current use of ≤3 drinks/week), moderate (current use of >3 to 7 drinks/week for women, and >3 to 14 drinks/week for men), and heavy (current use of >7 drinks/week for women and >14 drinks/week for men). In 1987 and 1992, the 3 abstainer categories could not be differentiated and were combined. In 1991, former and lifetime infrequent drinkers were combined but could be distinguished from never drinkers.

Mortality ascertainment. In 2004, the NCHS conducted an updated linkage of the NHIS to death certificate data found in the National Death Index. Linked mortality files include NHIS surveys from 1986 to 2000 with mortality follow-up from the month of interview through December 31, 2002. Mortality information is based upon the results

Abbreviations and Acronyms

- CHD** = coronary heart disease
- CI** = confidence interval
- CVD** = cardiovascular disease
- HDL-C** = high-density lipoprotein cholesterol
- HR** = hazard ratio
- NCHS** = National Center for Health Statistics
- NHIS** = National Health Interview Survey

Table 1 Alcohol-Related Variables Assessed in 9 Administrations of the NHIS

Year	n	CVD Deaths	How Assessed	Time Frame	Participants	Former Drinking	Binge Drinking	Adult Response Rate
1987	21,396	1,865	FFQ	Past yr	All	No	No	86%
1988	42,144	3,478	QF	Past yr	All	Yes	Yes	87%
1990	29,527	2,352	QF	Past 2 weeks*	All	Yes	No	83%
1991	19,956	85	QF	Past yr	Age 18–44 yrs	Yes†	Yes	75%
1992	11,076	631	FFQ	Past yr	All	No	No	86%
1997‡	33,477	888	QF	Past yr	All	Yes	Yes	80%
1998	29,761	639	QF	Past yr	All	Yes	Yes	74%
1999	28,206	427	QF	Past yr	All	Yes	Yes	70%
2000	29,664	305	QF	Past yr	All	Yes	Yes	72%

*Only individuals who reported that their use in the past 2 weeks was typical of their use in the past year were included in these analyses. †Former and lifetime infrequent drinking were combined in 1991. ‡The National Health Interview Survey (NHIS) administered a similar questionnaire from 1997 through 2000.
 CVD = cardiovascular disease; FFQ = food frequency questionnaire; QF = quantity-frequency.

from a probabilistic match between NHIS and National Death Index death certificate records (9). All NHIS participants 18 years and older were eligible for mortality follow-up. To minimize potential breaches of confidentiality, the NCHS developed a minor data perturbation plan; empirical studies suggest that the original and perturbed data files yield very similar descriptive statistics and model-based associations with mortality (10).

Deaths were coded for underlying cause with NCHS bridge codes spanning the Ninth and Tenth Revisions of the International Classification of Diseases (ICD-9, years 1979 to 1998; ICD-10, years 1999 to present) (11). We separately examined cardiovascular mortality (bridge codes 053 and 075, corresponding to ICD-10, I00 to I99), CHD mortality (bridge code 058, ICD-10, I20 to I25), and cerebrovascular mortality (bridge code 070, ICD-10, I60 to I69).

Covariates. We included a core set of covariates available in all NHIS surveys that also assessed alcohol consumption. We categorized self-reported race/ethnicity as non-Hispanic white, non-Hispanic black, Hispanic, and other. We grouped educational attainment as less than high school, high school, and some college, and marital status as married or nonmarried. We categorized region of residence in 4 U.S. Census regions (Northeast, South, Midwest, and West). Urbanization was categorized as residence in a U.S. census-defined metropolitan statistical area with a population of 1 million or more, a population of <1 million, or outside of a defined metropolitan statistical area. Smoking status was defined in 5 categories: never smokers, former smokers, and tertiles of intensity among current smokers. We calculated body mass index on the basis of self-reported height and weight and used standard World Health Organization categories of underweight, normal weight, overweight, and obese. Self-reported general health status was categorized as excellent, very good, good, and fair/poor/unknown.

The NCHS queried physical activity and diet only in selected NHIS administrations. Leisure-time physical activity was assessed with a series of items on specific activities in 1990 and 1991 (12,13) and general items about moderate and heavy physical exertion in 1997 to 2000 (14); to combine these, activity was categorized in quartiles. Diet was assessed with a food-frequency questionnaire in 1987 and 1992. We used, as previously described (15), foods grouped as fruits and vegetables, meat/poultry/fish, dairy products, added fats, and desserts, all categorized in tertiles.

Statistical analysis. We performed separate, similar analyses of alcohol consumption and mortality for each NHIS survey year with Cox proportional hazards regression, with appropriate sample weights provided by the NCHS. We set age as the metamer (16) and adjusted for sex and race in initial models and additionally for marital status, education, region, urbanization, smoking, body mass index, and general health status in multivariable models.

From the hazards ratios estimated in individual Cox proportional hazards models, we derived summary estimates of relative risk with random effects meta-analysis to combine across survey years (17,18). Because each survey provides estimates for the same underlying population (the noninstitutionalized adult population of the U.S.), we also performed pooled analyses in sensitivity analyses.

For primary analyses, we set never drinkers as a referent; because this excluded data from 1987 and 1992, we included these years in additional analyses with all nondrinkers as a referent only after comparing the risk associated with former and lifetime infrequent drinking. We tested several domains of alcohol consumption, including average volume of intake, frequency and quantity of intake, and binge drinking.

We performed sensitivity analyses with additional adjustment for diet or physical activity where such data were available, with exclusion of individuals who died within the first 2 years (i.e., a 2-year lag), and restricted to 5 years of follow-up for each survey (to reduce misclassification over longer follow-up). We performed pre-specified analyses stratified by sex, age (<55 years vs. 55 years and older), race (white vs. nonwhite), and general health (excellent vs. other); to maximize precision, we used all available NHIS years in stratified analyses. Given the possibility that low folate intake attenuates the lower risk of CHD associated with alcohol intake (19), we also examined risk stratified by follow-up that occurred before and after the initiation of folate fortification in the U.S. in 1998 (20).

Results

A total of 245,207 participants were included in the 9 iterations of the NHIS included in these analyses (Table 1), generally ranging from 20,000 to 40,000 participants/year. There were a total of 10,670 cardiovascular deaths during 1,987,439 person-years of follow-up, including 6,135 related to CHD and 1,758 related to stroke. Baseline characteristics according to alcohol consumption in the pooled sample are shown in the Online Tables.

Average volume of alcohol consumption. Figure 1 shows adjusted hazard ratios (HRs) for CVD mortality according to light, moderate, and heavy drinking at baseline in the individual NHIS surveys. In general, light and moderate drinking were associated with lower risk than abstention, with somewhat lower risk estimates for moderate drinking, whereas heavy drinking was not clearly associated with higher or lower risk.

Table 2 shows summary HRs for CVD mortality according to alcohol intake across 7 NHIS years with information on lifetime drinking habits. Risk tended to be similar among never drinkers and lifetime infrequent drinkers. A modestly higher risk was observed among former drinkers only in initial analyses and was eliminated by multivariable adjustment. In contrast, multivariable adjustment had relatively little effect upon risk estimates among current drinkers. There was no higher adjusted risk among former drinkers




Figure 1 Cardiovascular Mortality According to Alcohol Intake and NHIS Year

Adjusted hazard ratios (HRs) are shown relative to abstainers from alcohol. **T-bars** indicate 95% confidence intervals (CIs). CVD = cardiovascular disease; NHIS = National Health Interview Survey.

than never drinkers for CHD mortality (adjusted HR: 1.00; 95% confidence interval [CI]: 0.89 to 1.11) or cerebrovascular mortality (adjusted HR: 1.06; 95% CI: 0.86 to 1.33).

Table 3 shows the association of average alcohol intake with CVD, coronary, and stroke mortality across all NHIS years, both overall and in pre-specified subgroups. In general, the association of light and, especially, moderate drinking with lower risk was observed to similar degrees across all subgroups, although estimates tended to be further below 1 for coronary than stroke mortality, particularly for heavy drinking. Light and moderate drinking were also inversely associated with cardiovascular deaths not coded as ischemic and cerebrovascular, with HRs of 0.73 (95% CI: 0.64 to 0.83) and 0.71 (95% CI: 0.59 to 0.85), respectively.

Hazard ratios tended to be numerically closer to 1 among minority populations, although light and moderate drinking were associated with significantly lower risk even among minority participants. This contrast was more pronounced for CHD mortality, because light and moderate drinking were associated with summary HRs of 0.74 (95% CI: 0.67 to 0.82) and 0.66 (95% CI: 0.55 to 0.78), respectively, among non-Hispanic whites; the corresponding HRs among members of minority populations were 0.85 (95% CI: 0.67 to 1.08) and 0.79 (95% CI: 0.56 to 1.10).

We performed several sensitivity analyses to test our findings (Table 2). Additional adjustment for physical activity and diet altered the observed summary estimates to a minimal degree. When a 2-year lag was incorporated to

Table 2 Adjusted Summary HRs (and 95% CIs) for Cardiovascular Mortality Among NHIS Participants in 1988, 1990, 1991, and 1997 to 2000

	Volume of Alcohol Consumption					
	Never Drinker	Lifelong Infrequent Drinker	Former Drinker	Light	Moderate	Heavy
CVD deaths*	2,754	1,538	1,700	1,131	697	354
Person-yrs	305,966	242,849	174,714	520,562	238,498	94,815
Age-, sex-, and race-adjusted	1.00	0.98 (0.91-1.05)	1.15 (1.06-1.24)	0.66 (0.56-0.77)	0.58 (0.47-0.72)	1.07 (0.94-1.22)
Multivariable-adjusted†	1.00	0.95 (0.88-1.02)	1.02 (0.94-1.11)	0.69 (0.59-0.82)	0.62 (0.50-0.77)	0.95 (0.82-1.10)
MV activity‡	1.00	0.96 (0.89-1.03)	1.02 (0.94-1.11)	0.71 (0.61-0.83)	0.65 (0.53-0.79)	0.96 (0.83-1.11)
Pooled MV HR§	1.00	0.96 (0.89-1.03)	1.02 (0.94-1.10)	0.71 (0.65-0.78)	0.70 (0.63-0.78)	0.95 (0.82-1.10)
2-yr lag	1.00	0.96 (0.89-1.04)	1.03 (0.94-1.13)	0.72 (0.62-0.84)	0.74 (0.63-0.86)	1.04 (0.90-1.21)

According to baseline volume of alcohol consumption. *Hazard ratios (HRs) derive from random-effects meta-analyses of National Health Interview Survey (NHIS) years 1988, 1990, 1991, and 1997 to 2000; 1987 and 1992 did not include information with which to identify former drinkers. †Multivariable-adjusted HRs adjusted for age, sex, race, smoking, marital status, education, region, urbanization, smoking, body mass index, and general health status. ‡Analyses adjusted for covariates in multivariable (MV) models and physical activity assessed in 1990, 1991, and 1997 to 2000. §Hazard ratios derive from pooled (rather than meta-analyzed) analyses of individual NHIS administrations. ||Lagged analyses exclude individuals who died within the first 2 years after administration of the respective NHIS. CI = confidence interval; CVD = cardiovascular disease; HR = hazard ratio.

Table 3 Adjusted Summary HRs (and 95% CIs) for Cardiovascular, Coronary Heart Disease, and Stroke Mortality Among NHIS Participants in 1987, 1988, 1990 to 1992, and 1997 to 2000

	Volume of Alcohol Consumption			
	Nondrinkers	Light	Moderate	Heavy
Cardiovascular deaths	7,400	1,804	936	530
Person-yr	856,659	677,069	310,062	143,649
Multivariable-adjusted*	1.00	0.76 (0.68–0.85)	0.67 (0.59–0.77)	0.89 (0.73–1.10)
Multivariable + activity/diet†	1.00	0.77 (0.69–0.85)	0.69 (0.61–0.80)	0.90 (0.73–1.10)
5-yr follow-up‡	1.00	0.71 (0.60–0.84)	0.65 (0.55–0.77)	0.83 (0.66–1.04)
Men	1.00	0.78 (0.69–0.90)	0.70 (0.61–0.81)	0.95 (0.76–1.19)
Women	1.00	0.74 (0.66–0.83)	0.70 (0.60–0.81)	0.88 (0.72–1.07)
White	1.00	0.75 (0.68–0.84)	0.67 (0.58–0.78)	0.90 (0.73–1.10)
Not white	1.00	0.80 (0.65–0.99)	0.76 (0.61–0.94)	0.92 (0.70–1.21)
Age <55 yrs	1.00	0.76 (0.60–0.97)	0.62 (0.45–0.86)	0.91 (0.60–1.40)
Age ≥55 yrs	1.00	0.77 (0.69–0.86)	0.71 (0.63–0.80)	0.89 (0.76–1.06)
Excellent health	1.00	0.70 (0.56–0.88)	0.69 (0.57–0.83)	0.92 (0.66–1.28)
Not excellent health	1.00	0.72 (0.64–0.80)	0.61 (0.52–0.72)	0.82 (0.65–1.03)
Coronary heart disease deaths	4,217	1,057	553	308
Coronary mortality	1.00	0.75 (0.66–0.84)	0.67 (0.57–0.79)	0.80 (0.61–1.05)
Stroke deaths	1,241	288	135	94
Stroke mortality	1.00	0.80 (0.61–1.05)	0.76 (0.58–0.99)	1.25 (0.92–1.70)

According to baseline volume of alcohol consumption. *Multivariable-adjusted HRs adjusted for age, sex, race, smoking, marital status, education, region, urbanization, smoking, body mass index, and general health status. †Analyses adjusted for covariates in multivariable models; physical activity assessed in 1990, 1991, and 1997 to 2000; and diet assessed in 1987 and 1992. ‡Truncated analyses limited follow-up to 5 years after administration of the respective NHIS.

Abbreviations as in Tables 1 and 2.

eliminate early deaths, the observed HRs were attenuated but remained statistically significant for light and moderate drinking. In contrast, restriction to 5 years of follow-up for each survey administration to minimize misclassification of alcohol intake over time tended to increase the lower risk associated with alcohol intake. Pooled analyses yielded roughly similar estimates, albeit with substantially tighter confidence intervals. In general, we found no material differences in CVD mortality associated with light or moderate drinking in analyses stratified by follow-up before or after folate fortification in the U.S. (data not shown).

Frequency and quantity of alcohol consumption. Figure 2 demonstrates the joint associations of drinking frequency and quantity with cardiovascular mortality compared with abstainers. There was not a simple clear pattern, although risk was consistently higher among those who consumed 3 compared with 2 drinks/drinking days. There was no consistent association of drinking frequency with risk. These general findings were similar in analyses of CHD rather than CVD mortality and among both men and women.

Binge drinking. There was little overall association of binge drinking with CVD mortality. In analyses of NHIS years 1988, 1991, and 1997 to 2000, the multivariable-adjusted HRs for CVD mortality (compared with long-term abstainers) were 0.65 (95% CI: 0.51 to 0.81) among drinkers who reported no binge drinking, 0.66 (95% CI: 0.46 to 0.95) among those who endorsed binge drinking <1 day/week, and 0.68 (95% CI: 0.44 to 1.06) among those who reported binge drinking ≥1 day/week, consistent with the lower overall risk associated with drinking. After taking

usual drinking habits into account, we found limited evidence that binge drinking might attenuate the lower risk of CHD mortality associated with light and moderate drinking among women. For example, among women who did not report binge drinking, light and moderate drinking together were associated with a summary HR for CHD mortality of 0.56 (95% CI: 0.41 to 0.76), compared with long-term abstinence; the corresponding HR among binge drinkers




Figure 2 Cardiovascular Mortality According to Quantity and Frequency of Alcohol Consumption

Adjusted summary hazard ratios (HRs) are shown relative to abstainers from alcohol. CVD = cardiovascular disease.

was 1.23 (95% CI: 0.42 to 3.61). However, relatively few women reported binge drinking, and the difference in risk between light/moderate drinkers according to binge drinking was not significant ($p = 0.17$). Power was even more limited to assess the risk of binge drinking specifically associated with stroke mortality.

Discussion

In this analysis of 9 nationally representative samples of the U.S. population, comprising over 10,000 cardiovascular deaths, light and moderate alcohol consumption was associated with lower cardiovascular mortality, whereas heavy alcohol consumption was not.

Great interest remains in whether the previously observed relationships of alcohol consumption with lower rates of CVD or mortality are biased by choice of referent category (21,22). Because former drinkers might have ceased drinking for health-related reasons, they have typically been found to have higher rates of mortality in previous studies (23,24), although studies of CVD have not always demonstrated this increase in risk (25). Even less clear is the appropriate treatment of rare drinkers, because critics have variously suggested that they be fully excluded from the referent category of long-term abstainers (5) or replace abstainers as the referent (26). Our data demonstrate that, at least for CVD mortality, risk is similar among lifelong abstainers, lifelong rare drinkers, and former drinkers and thus that the inverse associations of light and moderate drinking with risk of CVD mortality are robust with respect to the choice of referent category.

Existing data on alcohol use and CVD derive nearly exclusively from studies of limited generalizability to the full U.S. population, although some might be representative of specific U.S. communities (27,28). As a result, our results provide some of the strongest evidence to date that the observed associations can be generalized to the U.S. population and are not limited to intensively monitored cohorts of volunteers. However, the NHIS excludes institutionalized individuals, who are likely to include the frailest members of society for whom even limited alcohol consumption could be detrimental.

Our results also provide insight into the consistency of the association of alcohol consumption and risk of CVD across subgroups. In general, light and moderate alcohol consumption were associated similarly with lower risk of CVD in nearly all segments of the U.S. population, including those who report themselves to be in excellent health (in whom the likelihood of residual confounding seems smallest) (29). One relative exception to this finding was race/ethnicity, where the association seemed stronger among non-Hispanic whites. Limited data suggest that the apparent U- or J-shaped associations of alcohol consumption with CVD and mortality observed in other groups might not extend to African Americans, who comprise the largest minority population in these surveys (30,31). Interestingly,

alcohol consumption was associated with a numerically higher risk of myocardial infarction among black Africans but a lower risk among white and mixed-race Africans in the Inter-Heart Int Africa study (32), suggesting that findings among African Americans might reflect known ancestral variation in alcohol dehydrogenase 1B isoforms or other alcohol-metabolizing enzymes (33). Prospective studies of alcohol consumption and CVD in minority populations are needed to clarify this issue.

The association of alcohol consumption with risk of CHD mortality tended to be stronger than that for cerebrovascular mortality in our analyses. This observation mirrors that observed for incident CVD in several other cohorts, where the associations of alcohol intake with CHD (27,34,35) have been stronger and more linear than the corresponding associations even with ischemic stroke (36–38). This might reflect, in part, the particularly strong contributions of hypertension and atrial fibrillation as risk factors for ischemic stroke, both of which are positively associated with at least heavy drinking (39,40). In the NHIS, the weaker association with cerebrovascular mortality likely also reflects some contribution of hemorrhagic stroke, which tends to be positively associated with alcohol consumption (41), perhaps related to the antiplatelet activity of alcohol (42).

Important limitations of our study warrant discussion. As with any observational study, our results might be confounded by factors for which we did not fully adjust, although the range of covariates available in the NHIS is relatively broad. Any unadjusted or incompletely unadjusted factor would need to be associated with both alcohol consumption and CVD and not strongly associated with the covariates already included in our models. Previous work in the Behavioral Risk Factor Surveillance System, another nationally representative survey, suggests that adjustment for race and education—which were included in these analyses—might be particularly important in reducing residual confounding (43).

The assessment of alcohol consumption in the NHIS occurred at a single point in time. This might have led us to underestimate the strength of associations for which alcohol consumption has acute effects on risk, although alcohol consumption is a relatively stable behavior among light- and moderate-drinking adults in the U.S. (44). It is most likely to have weakened associations of CVD with drinking frequency, where the presumed mechanisms of action include short-term, completely reversible effects on factors like thrombotic potential.

Although all data were collected prospectively with standardized, well-recognized instruments, the NHIS does not include measurement of biological factors that could be used to validate self-reported alcohol consumption (e.g., HDL-C) (45) or examine potential pathways that mediate the observed relationship (e.g., adiponectin) (46). Information on individual beverage types was also not uniformly collected, although most evidence suggests that wine, beer,

and spirits are similar in their relationships with HDL-C (47) and with risk of CVD (48).

Despite the large overall sample size of our analyses, we were nonetheless limited in power in some cases, such as when stratifying participants by binge drinking and usual drinking habits simultaneously. In part, this reflects the relative rarity of binge drinking among otherwise light-drinking women, but it nonetheless limits our ability to estimate the risk associated with binge drinking with precision. Similarly, power was insufficient to examine ischemic and hemorrhagic strokes separately; the overall association observed here likely reflects lower risk of more common ischemic strokes and higher risk of hemorrhagic strokes (49).

The NHIS obtains mortality data from the National Death Index, which derives cause-of-death information from state vital statistics offices and ultimately from death certificates. Although this methodology matches formal nosologist evaluations reasonably well (50), death certificates are nonetheless prone to misclassify CVD as a cause of death (51). Because this type of misclassification is likely random with respect to alcohol consumption, the true strength of the associations we evaluated is apt to be greater than that observed.

In summary, in aggregate analyses of 9 nationally representative surveys of the U.S. population conducted between 1987 and 2000, light and moderate alcohol consumption were inversely associated with cardiovascular mortality, with little difference in risk among lifelong abstainers, lifelong rare drinkers, and former drinkers. The magnitude of lower risk was generally strongest for CHD mortality and among non-Hispanic whites. These data bolster previous epidemiological studies that have found lower rates of incident CVD among moderate drinkers but also provide cautionary evidence that drinking above recommended limits eliminates this risk reduction. Although randomized trials to evaluate these relationships might yet be conducted to test these issues definitively, clinicians and patients at this time must continue to make informed, individualized, and collaborative decisions about the safety of alcohol consumption.

Acknowledgment

The authors thank Barry I. Graubard, PhD, of the National Cancer Institute's Division of Cancer Epidemiology and Genetics, for invaluable statistical advice.

Reprint requests and correspondence: Dr. Kenneth J. Mukamal, BIDMC Division of General Medicine and Primary Care Research Program, 1309 Beacon Street, 2nd Floor, Brookline, Massachusetts 02446. E-mail: kmukamal@bidmc.harvard.edu.

REFERENCES

1. Cleophas TJ. Wine, beer and spirits and the risk of myocardial infarction: a systematic review. *Biomed Pharmacother* 1999;53:417–23.
2. Di Castelnuovo A, Rotondo S, Iacoviello L, Donati MB, De Gaetano G. Meta-analysis of wine and beer consumption in relation to vascular risk. *Circulation* 2002;105:2836–44.
3. Rimm EB, Williams P, Fosher K, Criqui M, Stampfer MJ. Moderate alcohol intake and lower risk of coronary heart disease: meta-analysis of effects on lipids and haemostatic factors. *BMJ* 1999;319:1523–8.
4. Mukamal K. Alcohol intake and noncoronary cardiovascular diseases. *Ann Epidemiol* 2007;17:S8–12.
5. Fillmore KM, Kerr WC, Stockwell T, Chikritzhs T, Bostrom A. Moderate alcohol use and reduced mortality risk: systematic error in prospective studies. *Addict Res Theory* 2006;14:101–32.
6. Rehm JT, Bondy SJ, Sempos CT, Vuong CV. Alcohol consumption and coronary heart disease morbidity and mortality. *Am J Epidemiol* 1997;146:495–501.
7. Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos CT. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addiction* 2003;98:1209–28.
8. Block G, Subar AF. Estimates of nutrient intake from a food frequency questionnaire: the 1987 National Health Interview Survey. *J Am Diet Assoc* 1992;92:969–77.
9. National Center for Health Statistics Office of Analysis and Epidemiology. The 1986–2000 National Health Interview Survey Linked Mortality Files: Matching Methodology. Hyattsville, MD: National Center for Health Statistics, 2005.
10. Lochner K, Hummer RA, Bartee S, Wheatcroft G, Cox C. The public-use National Health Interview Survey linked mortality files: methods of reidentification risk avoidance and comparative analysis. *Am J Epidemiol* 2008;168:336–44.
11. Anderson RN, Minino AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: preliminary estimates. *Natl Vital Stat Rep* 2001;49:1–32.
12. Ainsworth B, Haskell W, Leon A, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc* 1993;25:71–80.
13. Prevalence of leisure-time and occupational physical activity among employed adults—United States, 1990. *MMWR Morb Mortal Wkly Rep* 2000;49:420–4.
14. Blackwell DL, Collins JG, Coles R. Summary health statistics for U.S. adults: National Health Interview Survey, 1997. *Vital Health Stat* 10 2002;1–109.
15. Breslow RA, Graubard BI, Sinha R, Subar AF. Diet and lung cancer mortality: a 1987 National Health Interview Survey cohort study. *Cancer Causes Control* 2000;11:419–31.
16. Korn EL, Graubard BI, Midthune D. Time-to-event analysis of longitudinal follow-up of a survey: choice of the time-scale. *Am J Epidemiol* 1997;145:72–80.
17. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
18. Rao RS, Graubard BI, Schmid CH, et al. Meta-analysis of survey data: application to health services research. *Health Serv Outcomes Res Methodol* 2008;8:98–114.
19. Rimm EB, Willett WC, Hu FB, et al. Folate and vitamin B6 from diet and supplements in relation to risk of coronary heart disease among women. *JAMA* 1998;279:359–64.
20. Choumenkovitch SF, Selhub J, Wilson PWF, Rader JI, Rosenberg IH, Jacques PF. Folic acid intake from fortification in United States exceeds predictions. *J Nutr* 2002;132:2792–8.
21. Rehm J, Irving H, Ye Y, Kerr WC, Bond J, Greenfield TK. Are lifetime abstainers the best control group in alcohol epidemiology? On the stability and validity of reported lifetime abstinence. *Am J Epidemiol* 2008;168:866–71.
22. Klatsky AL. Invited commentary: never, or hardly ever? It could make a difference. *Am J Epidemiol* 2008;168:872–5, discussion 876–7.
23. Di Castelnuovo A, Costanzo S, Bagnardi V, Donati MB, Iacoviello L, de Gaetano G. Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies. *Arch Intern Med* 2006;166:2437–45.
24. Gmel G, Gutjahr E, Rehm J. How stable is the risk curve between alcohol and all-cause mortality and what factors influence the shape? A precision-weighted hierarchical meta-analysis. *Eur J Epidemiol* 2003;18:631–42.
25. Jackson R, Scragg R, Beaglehole R. Alcohol consumption and risk of coronary heart disease. *BMJ* 1991;303:211–6.
26. Shaper AG. Alcohol and mortality: a review of prospective studies. *Br J Addict* 1990;85:837–47, discussion 849–61.

27. Mukamal KJ, Chung H, Jenny NS, et al. Alcohol consumption and risk of coronary heart disease in older adults: the Cardiovascular Health Study. *J Am Geriatr Soc* 2006;54:30–7.
28. Friedman LA, Kimball AW. Coronary heart disease mortality and alcohol consumption in Framingham. *Am J Epidemiol* 1986;124:481–9.
29. Mukamal KJ, Chiuve SE, Rimm EB. Alcohol consumption and risk for coronary heart disease in men with healthy lifestyles. *Arch Intern Med* 2006;166:2145–50.
30. Pletcher MJ, Varosy P, Kiefe CI, Lewis CE, Sidney S, Hulley SB. Alcohol consumption, binge drinking, and early coronary calcification: findings from the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Am J Epidemiol* 2005;161:423–33.
31. Sempos CT, Rehm J, Wu T, Crespo CJ, Trevisan M. Average volume of alcohol consumption and all-cause mortality in African Americans: the NHEFS cohort. *Alcohol Clin Exp Res* 2003;27:88–92.
32. Steyn K, Sliwa K, Hawken S, et al. Risk factors associated with myocardial infarction in Africa: the INTERHEART Africa study. *Circulation* 2005;112:3554–61.
33. Edenberg HJ, Xuei X, Chen HJ, et al. Association of alcohol dehydrogenase genes with alcohol dependence: a comprehensive analysis. *Hum Mol Genet* 2006;15:1539–49.
34. Klatsky AL, Armstrong MA, Friedman GD. Relations of alcoholic beverage use to subsequent coronary artery disease hospitalization. *Am J Cardiol* 1986;58:710–4.
35. Rimm EB, Giovannucci EL, Willett WC, et al. Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet* 1991;338:464–8.
36. Klatsky AL, Armstrong MA, Friedman GD, Sidney S. Alcohol drinking and risk of hospitalization for ischemic stroke. *Am J Cardiol* 2001;88:703–6.
37. Mukamal KJ, Chung H, Jenny NS, et al. Alcohol use and risk of ischemic stroke among older adults: the cardiovascular health study. *Stroke* 2005;36:1830–4.
38. Mukamal KJ, Ascherio A, Mittleman MA, et al. Alcohol and risk for ischemic stroke in men: the role of drinking patterns and usual beverage. *Ann Intern Med* 2005;142:11–9.
39. Klatsky AL. Alcohol and hypertension. *Clin Chim Acta* 1996;246:91–105.
40. Djousse L, Levy D, Benjamin EJ, et al. Long-term alcohol consumption and the risk of atrial fibrillation in the Framingham Study. *Am J Cardiol* 2004;93:710–3.
41. Reynolds K, Lewis B, Nolen JD, Kinney GL, Sathya B, He J. Alcohol consumption and risk of stroke: a meta-analysis. *JAMA* 2003;289:579–88.
42. Renaud SC, Ruf JC. Effects of alcohol on platelet functions. *Clin Chim Acta* 1996;246:77–89.
43. Mukamal KJ, Ding EL, Djousse L. Alcohol consumption, physical activity, and chronic disease risk factors: a population-based cross-sectional survey. *BMC Public Health* 2006;6:118.
44. Kerr WC, Fillmore KM, Bostrom A. Stability of alcohol consumption over time: evidence from three longitudinal surveys from the United States. *J Stud Alcohol* 2002;63:325–33.
45. Giovannucci E, Colditz G, Stampfer MJ, et al. The assessment of alcohol consumption by a simple self-administered questionnaire. *Am J Epidemiol* 1991;133:810–7.
46. Beulens JW, van Loon LJ, Kok FJ, et al. The effect of moderate alcohol consumption on adiponectin oligomers and muscle oxidative capacity: a human intervention study. *Diabetologia* 2007;50:1388–92.
47. van der Gaag MS, van Tol A, Vermunt SH, Scheek LM, Schaafsma G, Hendriks HF. Alcohol consumption stimulates early steps in reverse cholesterol transport. *J Lipid Res* 2001;42:2077–83.
48. Rimm EB, Klatsky A, Grobbee D, Stampfer MJ. Review of moderate alcohol consumption and reduced risk of coronary heart disease: is the effect due to beer, wine, or spirits. *BMJ* 1996;312:731–6.
49. Camargo CA Jr. Case-control and cohort studies of moderate alcohol consumption and stroke. *Clin Chim Acta* 1996;246:107–19.
50. Doody MM, Hayes HM, Bilgrad R. Comparability of national death index plus and standard procedures for determining causes of death in epidemiologic studies. *Ann Epidemiol* 2001;11:46–50.
51. Lloyd-Jones DM, Martin DO, Larson MG, Levy D. Accuracy of death certificates for coding coronary heart disease as the cause of death. *Ann Intern Med* 1998;129:1020–6.

Key Words: alcohol ■ epidemiology ■ mortality.

 **APPENDIX**

For supplementary tables, please see the online version of this article.